





<section-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header><section-header><text><list-item><list-item>

































## Treatment Highlights

• 6-mm Bar & 1-mm Wedge Wire Screens

O Activated Sludge

- Single Stage Nitrification Prior to 2017
- O Fine Bubble Aeration PanelsO Centrifugal & PD Blowers
- Airlift RAS Pumps
- O Ultraviolet Disinfection
- Autothermal Thermophilic Aerobic Digestion (ATAD)
- Centrifuge Dewatering









## 16/07/2019



















Microbial Analysis			
Sample Source	March 2018	April 2018	• Typical nitrifier population in
Sampling Method	ZeeLung cord		MLSS is 6-8%.
Quantification	qPCR <sup>(1)</sup>	RT-qPCR <sup>(2)</sup>	<ul> <li>ZeeLung biofilm is dominated by nitrifiers due to the counter- diffusion of O<sub>2</sub> in the biofilm.</li> </ul>
Method Total Nitrifier	40.5% ± 12.2%	29.5% ± 23.5%	
Population			
<sup>(1)</sup> amoA as primer for AOB; 16s rRNA primers for NOB <sup>(2)</sup> Quantitative reverse transcription PCR: RNA is transcribed into DNA, which is then used for qPCR analysis			
• Abundance of nitrifiers on the ZeeLung when compared to MLSS			





